

A reply to the criticism of our work (arXiv:1210.5501) by the DAMA collaboration

Josef Pradler*

*Department of Physics and Astronomy,
Johns Hopkins University, Baltimore, MD 21218, USA*

Itay Yavin†

*Perimeter Institute for Theoretical Physics 31 Caroline St. N,
Waterloo, Ontario, Canada N2L 2Y5 and
Department of Physics & Astronomy,
McMaster University 1280 Main St. W,
Hamilton, Ontario, Canada, L8S 4L8*

Abstract

We reply to a critique of our paper arXiv:1210.5501 by the DAMA collaboration which appeared in arXiv:1210.6199. Our original claim that the observed background levels are likely to require a large modulation fraction of any putative signal holds. In fact, in light of DAMA's recent comment our claim is further corroborated. We identify the source of the discrepancy between our own analysis and DAMA's claimed levels of unmodulated background. Our analysis indicates that the background in the signal region as reported by DAMA is likely underestimated.

* jpradler@pha.jhu.edu

† iyavin@perimeterinstitute.ca

In a recent publication [1] we pointed out that the empirical verification of a certain special decay branch of ^{40}K remains outstanding and discussed the importance of such a measurement. In addition, we discussed the general role ^{40}K plays as an important background in the DAMA experiment [2]. Shortly after our paper appeared, the DAMA collaboration has criticized some of its findings [3]. There are three points of substance in their critique:

Critique 1: The collaboration states that the contribution of this special decay of ^{40}K to the total ^{40}K contribution is only about 10%. They therefore criticize our discussion of this branch as "captious".

Critique 2: The collaboration states that the content of $^{\text{nat}}\text{K}$ has been measured by investigation of the double coincidence, is independent of the branching ratio into the ground state, and its average value over all crystals is 13 ppb. Statements based on 20 ppb are therefore wrong.

Critique 3: Finally, the collaboration claims an upper limit of 0.25 cpd/kg/keV for the unmodulated part of the signal in the 2-4 keV energy bin. Given a residual rate of 0.0194 ± 0.0022 cpd/kg/keV [2], this would allow for a modulation fraction of 6-10% which can be accommodated with many models of Dark Matter [4], contrary to our conclusions.

The first two issues are rather trivial. The third issue is very important and its elucidation requires understanding the details, which we give in the appendix. We will show that the upper limit of $S_0 \lesssim 0.25$ cpd/kg/keV is likely the result of an underestimation of the background in the signal region.

Reply 1: Independently of DAMA, the measurement of the ^{40}K decay directly to the ground state of ^{40}Ar is a missing piece in the experimental verification of leading-order predictions of weak decays, being the only such EC decay known of its kind (third unique-forbidden). In [1] we calculated the theoretical prediction for the branching ratio and motivated its measurement. The total decay of ^{40}K is a serious background in the DAMA experiment and it was in fact one of our findings that this special branch is subdominant to the one where ^{40}K decays into the ^{40}Ar excited state, followed by a

1.46 MeV gamma-ray which escapes detection. In their comment [3] DAMA quotes (for the first time in print) that the former contribution is 10% of the total ^{40}K contribution. This is indeed in good agreement with what we found using the theoretical branching ratio and using the independent Monte Carlo simulation of Ref. [5]. This agreement gives further credence to our treatment of potassium decays in DAMA.

Reply 2: Our results are presented for a whole range of $^{\text{nat}}\text{K}$ contaminations between 1-100 ppb including the DAMA quoted upper limit of 20 ppb, which appeared in the official publication describing the DAMA apparatus [6]. The collaboration now provides an average contamination level of 13 ppb in their comment [3]. We could not find this number in any of their published works but it can be found on slide number 8 in a talk by Nozzoli [7]. More importantly, no data is presented to support this number, nor does the collaboration provide the uncertainty associated with it. Nevertheless using 13 ppb we find that the required signal modulation fraction is above 20%. In [1] we identified a way to measure the contamination level in a manner that is free from the Monte Carlo modeling required by the coincidence method. Considering the immense importance of the precise amount of potassium background to any interpretation in terms of a dark matter signal the lack of details so far provided by the collaboration is unsettling.

Reply 3: Again, we could not find any mention of an upper limit of 0.25 cpd/kg/keV for the unmodulated signal part in any of DAMA's published works. We therefore have to resort to slide 8 of a talk by Nozzoli [7] where the limit is quoted and a curve for the background model is reported. In their comment, DAMA says that they account for this limit in [4] when assessing the viability of various DM scenarios. In Fig. 1 we show our reproduction of this background, which is a Gaussian centered at 3.2 keV together with a linear fit as explained in the Appendix. As is clear from the plot, the linearly rising function is not supported by the data above 4 keV (the statistical error bars are shown but are barely visible.) More importantly, the decrease towards lower energies strongly supports the notion that the background present in the signal region 2-4 keV is underestimated. As argued in our original work [1], a flat background component is expected because it is a universal feature of β^- decays for small electron velocities.

Indeed, adding a flat component greatly improves the fit as shown in Fig. 1. This results in a significantly larger background estimate in the signal region (see the Appendix for details). In light of this alternative estimate, the required modulation fractions quoted in our work [1], where a relatively low constant value of 0.85 cpd/kg/day was used, are in fact seen to be conservative.

What should be clear from this reply is that a more detailed and careful discussion of backgrounds by the DAMA collaboration is called for. That was among the main conclusions in [1] where we also identified measures which could help to clarify some of the uncertainties. We invite the interested readers to consult the appendix where more details are presented and to reproduce our results themselves.




FIG. 1. The dots (with error-bars) are the single-hit rate reported by DAMA in [6]. The thick red curve shows our reproduction of the background curve presented by DAMA in Ref. [7]. The dashed curve from 6.2 keV to 10 keV is the continuation of the linear trend. The linear trend is obtained by fitting the data between 5.3 keV and 10 keV and results in excellent agreement with the reported background [7], as explained in the Appendix. Using the same Gaussian, but with a better model for the background above 5 keV (flat + linear), we obtain the solid green curve. In blue, we show the resulting curve in the case when $^{\text{nat}}\text{K}$ contamination level is increased to 14 ppb instead. We emphasize that we did not fit to the data below 5 keV, but used the Gaussian as reported by the collaboration in [7].

Appendix A: Details of the analysis

The unmodulated background presented on slide 8 of Nozzoli [7] is reproduced accurately by fitting a linear function to the data points between 5.3 keV and 10 keV and adding to it a single Gaussian whose parameters are taken directly from ref. [7],

$$\text{BKG}_{\text{Nozzoli}}(E) = \frac{A}{\sqrt{2\pi}\sigma^2} \exp\left(-\frac{(E - E_{\text{K40}})^2}{2\sigma^2}\right) + \text{slope} \times E + \text{intercept}. \quad (\text{A-1})$$

The amplitude of the Gaussian is $A = 0.64$ which is in good agreement with the results of [5] with a $^{\text{nat}}\text{K}$ contamination of 13 ppb. The center and spread of the Gaussian are in good agreement with the expected 3.2 keV energy deposition ($E_{\text{K40}} = 3.15$ keV) and resolution ($\sigma = 0.618$ keV). One would naively expect that the quoted upper limit $S_0 \lesssim 0.25$ cpd/kg/keV was derived by subtracting this simple background model from the data. However, in contradiction to what the DAMA collaboration is quoting this yields only $S_0 \approx 0.14$ cpd/kg/keV in the 2-4 keV energy bin as can be easily checked.

Given the negligible statistical error bars, it is seen in Fig. 1 that the linear fit in Eq. (A-1) provides an inadequate description of the data between 5-10 keV where little DM signal is expected. A flat background up to about 7 keV where it is broken and followed by a linearly rising one provides a much better agreement with the data¹,

$$\text{BKG}_{\text{linear+flat}}(E) = \begin{cases} 0.866 & E < 7.05 \text{ keV} \\ 0.491 + 0.053E & E > 7.05 \text{ keV} \end{cases} \quad (\text{A-2})$$

Such a model is also physically well-motivated. The background from β^- -emitters is entirely flat at low energies. The rise above 7 keV is more difficult to interpret because the DAMA collaboration has not released the the immediate spectrum above 10 keV. For example, it could be a broad Iodine escape peak between 14-18 keV originating from external ^{210}Pb contamination. Though such features have been measured previously with NaI(Tl) crystals [8], without further insight into the spectrum this remains a speculation.

¹ We have also tried a more general model involving two linear fits, but the results are qualitatively unchanged - the fit between 5-7 keV prefers to be flat.




FIG. 2. The solid-red curve shows the background model presented in ref. [7]. The dashed-red curve is the extrapolation of the linear function as explained in the caption of Fig. 1. The solid-blue line represents a full fit to the unmodulated DAMA data composed of a Gaussian component, a flat component, and a linearly rising component. The center of the Gaussian, its spread, the flat piece, and the slope and intercept of the linear function were allowed to float (total of 5 parameters). We stress that the amplitude of the Gaussian was fixed to a level which corresponds to a 13 ppb of $^{\text{nat}}\text{K}$ contamination as quoted by the collaboration.

We have also verified that a more general fit to the entire data (including the width and center of the Gaussian) does not result in any substantial change to the flat component which becomes 0.86 cpd/kg/keV, in agreement with what we used in our original work [1]. In fact, this model results in a disturbingly good fit to the entire data set as shown in Fig. 2. We emphasize that this fit is done with a fixed amplitude for the Gaussian determined by a contamination level of $^{\text{nat}}\text{K}$ of 13 ppb as quoted by the collaboration. Only five parameters were allowed to float in the fit: the centre of the Gaussian; the resolution; the flat component; and the slope and intercept of linear trend. This surprising result shows how volatile the DM interpretation is to small changes in the background model and it again emphasizes the need for a more thorough discussion of backgrounds by the DAMA collaboration. In generating Fig. 1 above we attempted to remain as accommodating as possible and used the Gaussian

curve quoted in [7] for modeling the bump and not the fit shown in Fig. 2.

We notice that the residual signal that would result from Fig. 1 is so small that it would require a modulation fraction in excess of 50%. This is higher than the more conservative value we quoted in our paper [1] of a little over 20%. The reason for this is that the Gaussian model we assumed in our paper was based on the resolution as given by DAMA [6], which tends to be a little broader than what is actually observed in the bump. This results in slightly less potassium background in the signal region of 2-4 keV. Therefore a larger residual is possible and the modulation fraction is correspondingly smaller.

Finally, we stress that the purpose of the background analysis we present is not to claim a full understanding of the backgrounds. Our goal, as it was in our paper, is to call attention to this important background and encourage the collaboration to present a full account of its detailed understanding. That being said, the remarkable agreement of our simple background model with the data must give pause to anyone who wishes to interpret the DAMA results in terms of a signal of dark matter.

-
- [1] J. Pradler, B. Singh, and I. Yavin. On an unverified nuclear decay and its role in the DAMA experiment. 2012.
 - [2] R. Bernabei et al. New results from DAMA/LIBRA. *Eur.Phys.J.*, C67:39–49, 2010.
 - [3] R. Bernabei, P. Belli, F. Cappella, V. Caracciolo, R. Cerulli, et al. Comment on ‘On an unverified nuclear decay and its role in the DAMA experiment’ (arXiv:1210.5501). 2012.
 - [4] R. Bernabei et al. First results from DAMA/LIBRA and the combined results with DAMA/NaI. *Eur.Phys.J.*, C56:333–355, 2008.
 - [5] V.A. Kudryavtsev, M. Robinson, and N.J.C. Spooner. The expected background spectrum in NaI dark matter detectors and the DAMA result. *Astropart.Phys.*, 33:91–96, 2010.
 - [6] R. Bernabei et al. The DAMA/LIBRA apparatus. *Nucl.Instrum.Meth.*, A592:297–315, 2008.
 - [7] Talk given by F. Nozzoli. 2009. <http://taup2009.lngs.infn.it/slides/jul3/nozzoli.pdf>.
 - [8] Susana Cebrian, J. Amare, J.M. Carmona, E. Garcia, I.G. Irastorza, et al. Status of the ANAIS experiment at Canfranc. *Nucl.Phys.Proc.Suppl.*, 114:111–115, 2003.